

Stacky Curves in Characteristic p

Andrew J. Kobin

`akobin@ucsc.edu`


Madison Moduli Weekend

September 27, 2020




Introduction

Common problem: all sorts of information is lost when we consider quotient objects and/or singular objects.




Introduction

Solution: Keep track of lost information using *orbifolds* (topological and intuitive) or *stacks* (algebraic and fancy).



Introduction


Solution: Keep track of lost information using *orbifolds* (topological and intuitive) or *stacks* (algebraic and fancy).



Example: For the plane curve $X : y^2 - x = 0$, stacks remember automorphisms like $(x, y) \leftrightarrow (x, -y)$ using groupoids

Introduction

Solution: Keep track of lost information using *orbifolds* (topological and intuitive) or *stacks* (algebraic and fancy).




Goal: Classify stacky curves (= orbifold curves) in char. p
 (First steps: “Artin–Schreier Root Stacks”, arXiv:1910.03146)

Complex Orbifolds

Definition


A **complex orbifold** is a topological space admitting an atlas $\{U_i\}$ where each $U_i \cong \mathbb{C}^n/G_i$ for a finite group G_i , satisfying compatibility conditions (think: manifold atlas but with extra info).



Algebraic Stacks

There's also a version of orbifold in algebraic geometry: an **algebraic stack**.


Important class of examples we will focus on are **Deligne–Mumford stacks** \approx smooth varieties or schemes with a finite automorphism group attached at each point.



Algebraic Stacks

There's also a version of orbifold in algebraic geometry: an **algebraic stack**.


Important class of examples we will focus on are **Deligne–Mumford stacks** \approx smooth varieties or schemes with a finite automorphism group attached at each point.



Algebraic Stacks

There's also a version of orbifold in algebraic geometry: an **algebraic stack**.


Important class of examples we will focus on are **Deligne–Mumford stacks** \approx smooth varieties or schemes with a finite automorphism group attached at each point.



Algebraic Stacks

There's also a version of orbifold in algebraic geometry: an **algebraic stack**.


Important class of examples we will focus on are **Deligne–Mumford stacks** \approx smooth varieties or schemes with a finite automorphism group attached at each point.



Algebraic Stacks

There's also a version of orbifold in algebraic geometry: an **algebraic stack**.


Important class of examples we will focus on are **Deligne–Mumford stacks** \approx smooth varieties or schemes with a finite automorphism group attached at each point.



Algebraic Stacks

There's also a version of orbifold in algebraic geometry: an **algebraic stack**.

Important class of examples we will focus on are **Deligne–Mumford stacks** \approx smooth varieties or schemes with a finite automorphism group attached at each point.




Focus on curves for the rest of the talk

An Example

Example

The (compactified) moduli space of complex elliptic curves is a stacky \mathbb{P}^1 with a generic $\mathbb{Z}/2$ and a special $\mathbb{Z}/4$ and $\mathbb{Z}/6$.



Consequence: can deduce dimension formulas for modular forms from Riemann–Roch formula for stacks.

Goal: Classify stacky curves in char. p .

Main obstacle to overcome:

- In char. 0, local structure is determined by a cyclic group action.
- In char. p , **this is not enough information** – need more invariants than just the order of a cyclic group.

Results (K. '20):

- Every p -cover of curves factors étale-locally through an **Artin–Schreier root stack**.
- Every stacky curve with order p automorphism group is étale-locally an Artin–Schreier root stack.
- For any algebraic curve X , there are infinitely many non-isomorphic Deligne–Mumford stacks with coarse space X and degree p automorphism groups at the same sets of points.

Root Stacks

Key fact: in char. 0, all stabilizers (automorphism groups) are *cyclic*.

So stacky curves can be locally modeled by a *root stack*: charts look like

$$U \cong [\mathrm{Spec} A / \mu_n]$$

where $A = K[y]/(y^n - \alpha)$ and μ_n is the group of n th roots of unity.

(Think: degree n branched cover mod μ_n -action, but remember the action using groupoids.)

Root Stacks

More rigorously:

Definition (Cadman '07, Abramovich–Olsson–Vistoli '08)

Let X be a scheme and $L \rightarrow X$ a line bundle with section $s : X \rightarrow L$. The **n th root stack** of X along (L, s) is the fibre product

$$\begin{array}{ccc} \sqrt[n]{(L, s)/X} & \longrightarrow & [\mathbb{A}^1/\mathbb{G}_m] & x \\ \downarrow & (L, s) & \downarrow & \downarrow \\ X & \longrightarrow & [\mathbb{A}^1/\mathbb{G}_m] & x^n \end{array}$$

Here, $[\mathbb{A}^1/\mathbb{G}_m]$ is the classifying stack for pairs (L, s) .

Interpretation: $\sqrt[n]{(L, s)/X}$ admits a canonical tensor n th root of (L, s) , i.e. (M, t) such that $M^{\otimes n} = L$ and $t^n = s$ (after pullback).

Root Stacks

Theorem (Geraschenko–Satriano '15)

Every smooth separated **tame** Deligne–Mumford stack of finite type with trivial generic stabilizer is* a root stack over its coarse space.

Corollary

Tame stacky curves are completely described by their coarse space and a finite list of numbers corresponding to the orders of cyclic stabilizers at a finite number of stacky points.



Root Stacks

Theorem (Geraschenko–Satriano '15)

Every smooth separated **tame** Deligne–Mumford stack of finite type with trivial generic stabilizer is* a root stack over its coarse space.

Corollary

Tame stacky curves are completely described by their coarse space and a finite list of numbers corresponding to the orders of cyclic stabilizers at a finite number of stacky points.



What happens with **wild** stacky curves in char. p ?

Artin–Schreier Root Stacks

In trying to classify **wild** stacky curves in char. p , we face the following problems:

- ① Stabilizer groups need not be cyclic (or even abelian)
- ② Cyclic $\mathbb{Z}/p^n\mathbb{Z}$ -covers of curves occur in families
- ③ Root stacks don't work
 - Finding $M^{\otimes p}$ is a problem
 - $[\mathbb{A}^1/\mathbb{G}_m] \rightarrow [\mathbb{A}^1/\mathbb{G}_m], x \mapsto x^p$ is a problem

Key case: cyclic $\mathbb{Z}/p\mathbb{Z}$ stabilizers

Artin–Schreier Root Stacks

Idea: replace **tame** cyclic covers $y^n = f(x)$ with **wild** cyclic covers $y^p - y = f(x)$.

More specifically: **Artin–Schreier theory** classifies cyclic degree p -covers of curves in terms of the **ramification jump** (e.g. if $f(x) = x^m$ then m is the jump).

This suggests introducing wild stacky structure using the local model

$$U = [\mathrm{Spec} A / (\mathbb{Z}/p)]$$

where $A = k[y]/(y^p - y - f(x))$ and \mathbb{Z}/p acts additively.

Artin–Schreier Root Stacks

How do we do it?

$$\begin{array}{ccc}
 \sqrt[n]{(L, s)/X} & \longrightarrow & [\mathbb{A}^1/\mathbb{G}_m] \\
 \downarrow & & \downarrow \\
 X & \xrightarrow{(L, s)} & [\mathbb{A}^1/\mathbb{G}_m]
 \end{array}
 \qquad
 \begin{array}{c}
 x \\
 \downarrow \\
 x^n
 \end{array}$$

Artin–Schreier Root Stacks

How do we do it?

$$\begin{array}{ccc}
 \sqrt[n]{(L, s)/X} & \longrightarrow & [\mathbb{P}^1/\mathbb{G}_a] \\
 \downarrow & & \downarrow \\
 X & \xrightarrow{(L, s)} & [\mathbb{P}^1/\mathbb{G}_a]
 \end{array}
 \qquad
 \begin{array}{c}
 [u, v] \\
 \downarrow \\
 [u^p, v^p - vu^{p-1}]
 \end{array}$$

Artin–Schreier Root Stacks

How do we do it?

$$\begin{array}{ccc}
 \sqrt[n]{(L, s)/X} & \longrightarrow & [\mathbb{P}^1/\mathbb{G}_a] \\
 \downarrow & & \downarrow \\
 X & \xrightarrow{(L, s, f)} & [\mathbb{P}^1/\mathbb{G}_a]
 \end{array}
 \qquad
 \begin{array}{c}
 [u, v] \\
 \downarrow \\
 [u^p, v^p - vu^{p-1}]
 \end{array}$$

Artin–Schreier Root Stacks

How do we do it?

$$\begin{array}{ccc}
 \varphi_1^{-1}((L, s, f)/X) & \longrightarrow & [\mathbb{P}^1/\mathbb{G}_a] \\
 \downarrow & & \downarrow \\
 X & \xrightarrow{(L, s, f)} & [\mathbb{P}^1/\mathbb{G}_a]
 \end{array}
 \qquad
 \begin{array}{c}
 [u, v] \\
 \downarrow \\
 [u^p, v^p - vu^{p-1}]
 \end{array}$$

Artin–Schreier Root Stacks

Definition (K.)

Fix $m \geq 1$. Let X be a scheme, $L \rightarrow X$ a line bundle and $s : X \rightarrow L$ and $f : X \rightarrow L^{\otimes m}$ two sections not vanishing simultaneously. The **Artin–Schreier root stack** of X with jump m along (L, s, f) is the normalized pullback

$$\begin{array}{ccc}
 \wp_m^{-1}((L, s, f)/X) & \longrightarrow & [\mathbb{P}(1, m)/\mathbb{G}_a] & & [u, v] \\
 \downarrow \nu & & \downarrow & & \downarrow \\
 X & \xrightarrow{(L, s, f)} & [\mathbb{P}(1, m)/\mathbb{G}_a] & & [u^p, v^p - vu^{m(p-1)}]
 \end{array}$$

where

- $\mathbb{P}(1, m)$ is the weighted projective line with weights $(1, m)$
- $\mathbb{G}_a = (k, +)$, acting additively
- $[\mathbb{P}(1, m)/\mathbb{G}_a]$ is the classifying stack for triples (L, s, f) up to the principal part of f .

Artin–Schreier Root Stacks

$$\begin{array}{ccc}
 \varphi_m^{-1}((L, s, f)/X) & \longrightarrow & [\mathbb{P}(1, m)/\mathbb{G}_a] \\
 \downarrow \nu & & \downarrow \\
 X & \xrightarrow{(L, s, f)} & [\mathbb{P}(1, m)/\mathbb{G}_a]
 \end{array}
 \quad
 \begin{array}{c}
 [u, v] \\
 \downarrow \\
 [u^p, v^p - vu^{m(p-1)}]
 \end{array}$$

Interpretation: $\varphi_m^{-1}((L, s, f)/X)$ admits a canonical p th root of L , i.e. a line bundle M such that $M^{\otimes p} = L$, and an AS root of s .

Artin–Schreier Root Stacks

Key example:

Example (K.)

Consider the AS cover

$$\begin{array}{c}
 Y : y^p - y = x^{-m} \\
 \mathbb{Z}/p \downarrow \\
 \mathbb{P}^1 = \text{Proj } k[x_0, x_1]
 \end{array}$$

where k is an algebraically closed field of characteristic p . Then

$$\varrho_m^{-1}((\mathcal{O}(1), x_0, x_1^m)/\mathbb{P}^1) \cong [Y/(\mathbb{Z}/p)].$$

In general, every AS root stack is étale-locally isomorphic to such an “elementary AS root stack”.

Classification of (Some) Wild Stacky Curves

So let's classify us some wild stacky curves!
 (Assume: everything defined over $k = \bar{k}$)

Theorem 1 (K. '20)

Every Galois cover of curves $\varphi : Y \rightarrow X$ with an inertia group \mathbb{Z}/p factors étale-locally through an Artin–Schreier root stack:

$$\begin{array}{ccc}
 Y & \xrightarrow{\varphi} & X \\
 \text{ét} \uparrow & & \text{ét} \uparrow \\
 V & \longrightarrow \wp_m^{-1}((L, s, f)/U) \longrightarrow & U
 \end{array}$$

Informal consequence: there are infinitely many non-isomorphic stacky curves over \mathbb{P}^1 with a single stacky point of order p .

This phenomenon only occurs in char. p .

Classification of (Some) Wild Stacky Curves

Main result:

Theorem 2 (K. '20)

Every stacky curve \mathcal{X} with a stacky point of order p is étale-locally isomorphic to an Artin–Schreier root stack $\wp_m^{-1}((L, s, f)/U)$ over an open subscheme U of the coarse space of \mathcal{X} .

This can even be done globally if \mathcal{X} has coarse space \mathbb{P}^1 :

Theorem 3 (K. '20)

If \mathcal{X} has coarse space \mathbb{P}^1 and all stacky points of \mathcal{X} have order p , then \mathcal{X} is isomorphic to a fibre product of AS root stacks of the form $\wp_m^{-1}((L, s, f)/\mathbb{P}^1)$ for $(m, p) = 1$ and (L, s, f) .

Generalizations

What about \mathbb{Z}/p^2 -covers, stacky points of order p^2 , and beyond?

For cyclic stabilizer groups \mathbb{Z}/p^n , Artin–Schreier theory is subsumed by **Artin–Schreier–Witt theory**:

- AS equations $y^p - y = f(x)$ are replaced by Witt vector equations $\underline{y}^p - \underline{y} = \underline{f}(\underline{x}) = (f_0(\underline{x}), \dots, f_n(\underline{x}))$.
- Covers are characterized by *sequences of ramification jumps*.
- Local structure is $U = [\mathrm{Spec} A / (\mathbb{Z}/p^n)]$ where

$$A = K[\underline{y}] / (\underline{y}^p - \underline{y} - \underline{f})$$

where $\underline{f} = (f_0, \dots, f_{n-1})$ is a Witt vector over \overline{K} .

This local structure can be formally introduced using **Artin–Schreier–Witt root stacks** (work in progress).

Generalizations

Thank you!